

# THE PROMPT GAMMA-RAY AND AFTERGLOW ENERGIES OF SHORT-DURATION GAMMA-RAY BURSTS

E. BERGER<sup>1,2,3</sup>

*Draft version February 5, 2008*

## ABSTRACT

I present an analysis of the  $\gamma$ -ray and afterglow energies of the complete sample of 17 short-duration GRBs with prompt X-ray follow-up. I find that 80% of the bursts exhibit a linear correlation between their  $\gamma$ -ray fluence and the afterglow X-ray flux normalized to  $t = 1$  d, a proxy for the kinetic energy of the blast wave ( $F_{X,1} \propto F_{\gamma}^{1.01 \pm 0.09}$ ). An even tighter correlation is evident between  $E_{\gamma, \text{iso}}$  and  $L_{X,1}$  for the subset of 13 bursts with measured or constrained redshifts. The remaining 20% of the bursts have values of  $F_{X,1}/F_{\gamma}$  that are suppressed by about three orders of magnitude, likely because of low circumburst densities (Nakar 2007). These results have several important implications: (i) The X-ray luminosity is generally a robust proxy for the blast wave kinetic energy, indicating  $\nu_X > \nu_c$  and hence a circumburst density  $n \gtrsim 0.05 \text{ cm}^{-3}$ ; (ii) most short GRBs have a narrow range of  $\gamma$ -ray efficiency, with  $\langle \epsilon_{\gamma} \rangle \approx 0.85$  and a spread of 0.14 dex; and (iii) the isotropic-equivalent energies span  $10^{48} - 10^{52}$  erg. Furthermore, I find tentative evidence for jet collimation in the two bursts with the highest  $E_{\gamma, \text{iso}}$ , perhaps indicative of the same inverse correlation that leads to a narrow distribution of true energies in long GRBs. I find no clear evidence for a relation between the overall energy release and host galaxy type, but a positive correlation with duration may be present, albeit with a large scatter. Finally, I note that the outlier fraction of 20% is similar to the proposed fraction of short GRBs from dynamically-formed neutron star binaries in globular clusters. This scenario may naturally explain the bimodality of the  $F_X/F_{\gamma}$  distribution and the low circumburst densities without invoking speculative kick velocities of several hundred  $\text{km s}^{-1}$ .

*Subject headings:* gamma-rays:bursts

## 1. INTRODUCTION

One of the fundamental parameters of any explosive phenomenon is the overall energy release. In the specific case of gamma-ray bursts (GRBs), it is essential to measure the energy of both the prompt  $\gamma$ -ray phase ( $E_{\gamma}$ ) and the blast wave which powers the afterglow ( $E_K$ ), since these two quantities define the total relativistic output of the central engine ( $E_{\text{rel}}$ ) and the efficiency of energy dissipation in the form of  $\gamma$ -rays ( $\epsilon_{\gamma} \equiv E_{\gamma}/E_{\text{rel}}$ ). While the isotropic-equivalent prompt energy is easily derived from the observed fluence ( $F_{\gamma}$ ) and distance ( $d_L(z)$ ) of the burst, a measurement of  $E_K$  is more challenging because it requires observations of the afterglow over a broad range in time and frequency (e.g., Berger et al. 2000; Panaitescu & Kumar 2002; Yost et al. 2003). Luckily, the afterglow flux at frequencies above the synchrotron cooling frequency,  $\nu_c$ , provides a robust proxy for  $E_K$  since it is independent of the circumburst density and depends only weakly on the fraction of shock energy in magnetic fields,  $\epsilon_B$  (Kumar 2000; Freedman & Waxman 2001; Berger et al. 2003a). For typical parameters (Panaitescu & Kumar 2002; Berger et al. 2003a), the soft X-ray band lies above  $\nu_c$ , and therefore the X-ray luminosity,  $L_X \propto \epsilon_e E_K$ ; here  $\epsilon_e$  is the fraction of shock energy in relativistic electrons.

In the case of the long-duration GRBs ( $T_{90} \gtrsim 2$  s), ex-

tensive studies of the relativistic energy output based on these various techniques suggest that for most bursts  $E_{\gamma}$ ,  $E_K$ , and most importantly their sum, span less than an order of magnitude around  $\sim 10^{51}$  erg *when corrected for jet collimation of the ejecta* (Frail et al. 2001; Panaitescu & Kumar 2002; Berger et al. 2003a; Bloom et al. 2003; Berger et al. 2003b). Recent observations of low redshift long GRBs paint a more complicated picture in which a population of sub-energetic bursts ( $\sim 10^{50}$  erg) likely dominates the overall GRB rate (Soderberg et al. 2004, 2006b). Still, the conclusion that the energy release of long GRBs is in the range typical of supernovae, and that they are collimated with opening angles of  $\theta_j \sim 5^\circ$ , has placed valuable constraints on the progenitor and engine models.

The discovery of afterglow emission from short-duration GRBs less than 2 years ago makes it now possible to carry out a similar analysis of their total energy output. Early observations indicated a lower isotropic-equivalent energy release compared to long GRBs,  $\sim 10^{48} - 10^{49}$  erg, in both the prompt and afterglow phase (Berger et al. 2005b; Fox et al. 2005; Gehrels et al. 2005; Bloom et al. 2006), as well as wider jet opening angles (Burrows et al. 2006; Soderberg et al. 2006a). In addition, Nakar (2007) noted based on a small sample of six bursts that the ratio of afterglow X-ray flux and  $\gamma$ -ray fluence,  $F_X t/F_{\gamma}$ , appears to have a bimodal distribution with values of  $\sim 10^{-2}$  and  $\lesssim 10^{-4}$ .

<sup>1</sup>Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101

<sup>2</sup>Princeton University Observatory, Peyton Hall, Ivy Lane, Princeton, NJ 08544

<sup>3</sup>Hubble Fellow

They argue that the latter bursts require a low circumburst density,  $n \lesssim 10^{-5} \text{ cm}^{-3}$ , likely due to ejection of the progenitor system from its host galaxy into the intergalactic medium<sup>4</sup>.

The recent association of several short GRBs with faint galaxies, of which several have been spectroscopically confirmed to be at  $z \sim 0.4 - 1.1$ , indicates a wide redshift distribution, and hence a wider range of energies than previously proposed (Berger et al. 2006). Here, I take advantage of these new observations to study the prompt and afterglow energies of the complete sample of 17 short bursts with prompt X-ray follow-up, a much larger sample than previously investigated. Of these 17 bursts, eight have measured redshifts, and an additional five have approximate or constrained redshifts. Using the  $\gamma$ -ray fluences and X-ray fluxes normalized to 1 day after the burst I show that  $E_{\gamma, \text{iso}}$  and  $E_{K, \text{iso}}$  span at least three orders of magnitude (possibly up to  $\sim 10^{52} \text{ erg}$ ). More importantly, I establish that 80% of the bursts exhibit a tight correlation between their prompt and afterglow energy output, suggesting a narrow distribution of  $\epsilon_\gamma$ , while 20% indeed have ratios of  $F_X/F_\gamma$  that are suppressed by three orders of magnitude. These results place important constraints on the progenitor and energy extraction models, as well as the properties of the circumburst medium and the shock microphysical parameters.

## 2. PROMPT GAMMA-RAY AND X-RAY AFTERGLOW ENERGIES

The  $\gamma$ -ray and X-ray afterglow properties of the 17 short bursts with prompt X-ray follow-up are listed in Table 1. I also list the available measurements or constraints on the redshift from host galaxy and afterglow observations. The relevant references are provided in the Table.

For GRB 061210 the X-ray fluxes have not been published and I therefore obtained the XRT data from the High Energy Astrophysics Science Archive Research Center<sup>5</sup>. I processed the data with the `xrtpipeline` script packaged within the HEASoft software, using the default grade selection and screening parameters to produce a light curve for the 0.3 – 10 keV energy range. Using a Galactic neutral hydrogen column density,  $N_H \approx 3.4 \times 10^{20} \text{ cm}^{-2}$ , and a typical  $\beta_X \approx -1.1$ , I find a conversion rate of 1 count  $\text{s}^{-1} = 4 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ . In addition, inspection of the X-ray light curve of GRB 061006 (Schady et al. 2006) reveals a possible steepening at  $t \sim 10^5 \text{ s}$ . To investigate this possibility I obtained the XRT data from HEASARC and followed the same analysis procedure as for GRB 061210. I find a likely steepening using data that were obtained 3 – 4 days after the burst, with a flux decay rate,  $\alpha_X \approx -1.9$ .

In the analysis below I use the standard cosmological parameters:  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_m = 0.27$ , and  $\Omega_\Lambda = 0.73$ .

In Figure 1, I plot the histograms of  $F_\gamma$  and  $F_{X,1}$ , the X-ray flux normalized to  $t = 1 \text{ d}$  after the burst. I use the observed values of  $\alpha_X$  when available, or the median value,  $\langle \alpha_X \rangle \approx -1.1$ , when the decay rate is not known. The ma-

jority of the bursts span about two orders of magnitude around  $F_{X,1} \sim 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ , similar to the width of the distribution for long GRBs (Berger et al. 2003a, 2005a). However, there is a noticeable tail extending to fainter fluxes, which leads to an overall spread of about five orders of magnitude and a median flux of  $3 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ . The fluence distribution is narrower, with an overall width of about 2.7 orders of magnitude around a median value of  $2 \times 10^{-7} \text{ erg cm}^{-1}$ . The median values of both  $F_\gamma$  and  $F_{X,1}$  are about a factor of five smaller than for *Swift* long GRBs (Berger et al. 2005a).

More interesting is the apparent correlation between  $F_\gamma$  and  $F_{X,1}$  for the majority of the bursts (Figure 1). The Spearman’s rank correlation coefficient is  $\rho = 0.86$ , indicating a null hypothesis (no correlation) probability of only  $5.3 \times 10^{-5}$ . Fitting the data, I find that the best-fit relation is linear,  $F_{X,1} = 10^{-13.47 \pm 0.06} \times (F_\gamma / 10^{-6.68})^{1.01 \pm 0.09} \text{ erg cm}^{-2} \text{ s}^{-1}$  with a scatter of about 0.4 dex based on a fit to 13 of the 17 bursts that follow the trend. The tight linear correlation suggests that  $\epsilon_\gamma$  has a narrow distribution, and that in most bursts the X-ray band indeed lies above the cooling frequency and traces  $E_K$ . Based on the observed correlation I infer that  $\epsilon_e E_K / E_\gamma \propto t_1 F_{X,1} / F_\gamma \approx 0.016$  (see also Nakar 2007). For a typical  $\epsilon_e \sim 0.1$ , the fraction of energy emitted in  $\gamma$ -rays is thus  $\epsilon_\gamma \sim 85\%$ , with an overall range of about 55 – 95%, similar to the values inferred for long GRBs (Panaitescu & Kumar 2002).

An even tighter correlation is evident between the X-ray luminosity,  $L_{X,1} \propto \epsilon_e E_{K, \text{iso}}$ , and  $E_{\gamma, \text{iso}}$ ; Figure 2. The isotropic X-ray luminosity is given by:

$$L_{X,1} = 4\pi d_L^2 F_{X,1} (1+z)^{\alpha_X - \beta_X - 1}, \quad (1)$$

where for  $\nu_X > \nu_c$ ,  $\alpha_X - \beta_X = (2 - p)/4$ ,  $p \sim 2 - 3$  is the power law index of the electron distribution,  $N(\gamma) \propto \gamma^{-p}$ , and  $d_L$  is the luminosity distance. The isotropic  $\gamma$ -ray energy (neglecting redshift-dependent bolometric corrections) is given by:

$$E_{\gamma, \text{iso}} = 4\pi d_L^2 F_\gamma (1+z)^{-1}. \quad (2)$$

To assess  $L_{X,1}$  and  $E_{\gamma, \text{iso}}$  I use the redshifts measured for eight bursts from their host galaxies, as well as approximate redshifts for three bursts (050813, 060121, and 061201) and limits for an additional two (051210 and 060313). The Spearman’s rank correlation coefficient is  $\rho = 0.94$ , (null hypothesis probability of only  $3.2 \times 10^{-6}$ ). The correlation appears to be slightly non-linear,  $L_{X,1} \propto E_{\gamma, \text{iso}}^{1.13 \pm 0.06}$ , primarily because for  $p > 2$  the dependence on  $(1+z)$  does not cancel out when taking the ratio of Equations 1 and 2. As can be seen from Table 1, the bolometric correction factors to  $E_{\gamma, \text{iso}}$  are in the range of  $\sim 3 - 15$  when converting from the 15 – 150 keV band to the 20 – 2000 keV band. These upward corrections to  $E_{\gamma, \text{iso}}$  will tend to linearize the correlation.

In addition to the overall strong correlation, it is also clear from Figure 2 that the range of inferred energies spans  $E_{\gamma, \text{iso}} \sim 10^{48} - 10^{51} \text{ erg}$  (15 – 150 keV), and may exceed  $\sim 10^{52} \text{ erg}$  if some of the bursts with unknown redshifts are located at  $z \gtrsim 1$ , as suggested by their faint host

<sup>4</sup>Nakar (2007) also propose that a wide dispersion in the shock parameters or  $\epsilon_\gamma$  may lead to the bimodal distribution.

<sup>5</sup><http://heasarc.gsfc.nasa.gov/W3Browse/>

galaxies ( $R \gtrsim 23$  mag; Berger et al. 2006). The full range of 3 – 4 orders of magnitude is similar to the spread in  $E_{\gamma,\text{iso}}$  observed for long GRBs (Frail et al. 2001; Bloom et al. 2003).

### 3. IMPLICATIONS

#### 3.1. A Narrow Distribution of $\epsilon_\gamma$

The data presented in this paper conclusively show that the majority (80%) of all short GRBs exhibit a tight correlation between their isotropic-equivalent prompt  $\gamma$ -ray and blast wave kinetic energies (see also Nakar 2007). The inferred  $\gamma$ -ray efficiency is  $\sim 85\%$  with a narrow spread of about 0.14 dex. This result is indeed verified by the three bursts for which detailed afterglow observations are available, with derived  $\epsilon_\gamma$  values of 0.8 (GRB 050709; Fox et al. 2005), 0.2 (GRB 050724; Berger et al. 2005b), and 0.65 (GRB 051221a; Soderberg et al. 2006a).

The remaining 20% of short bursts have X-ray fluxes that are suppressed by about three orders of magnitude compared to their  $\gamma$ -ray fluence. I stress that if this is indeed due to a low circumburst density (§3.4), which leads to  $\nu_X < \nu_c$ , then observations of the afterglow at higher X-ray energies (above  $h\nu_c$ ) should recover the same narrow distribution of  $\epsilon_\gamma$  seen for the bulk of the population.

The observed correlation and the inferred narrow distribution of  $\epsilon_\gamma$  have several crucial implications for short GRB progenitor models, the energy extraction mechanism, and burst properties such as the circumburst density and shock microphysics. The overall narrow spread in  $\epsilon_\gamma$ , and the median value, are similar to those observed in long GRBs (Panaitescu & Kumar 2002), suggesting that the properties of the relativistic outflow are similar for long and short bursts and are generally independent of the identity of the progenitor or the circumburst environment.

In addition, the tight correlation between  $\gamma$ -ray and afterglow energies indicates that for most bursts the underlying assumption that  $\nu_X > \nu_c$  is indeed correct. Therefore the required circumburst densities are (Granot & Sari 2002)  $n \gtrsim 0.05 (1+z)^{-1/2} \epsilon_{B,-1}^{-3/2} E_{50}^{1/2} \text{ cm}^{-3}$ , typical of interstellar environments. In the context of binary compact object progenitors (NS-NS, NS-BH), the inferred densities indicate that the majority of the progenitors do not experience kick velocities that are large enough for ejection into the intergalactic medium and/or that the merger timescales are short enough that the distance traveled is  $\lesssim 10$  kpc. Finally, the fraction of energy in relativistic electrons,  $\epsilon_e \approx (1 - \epsilon_\gamma) E_K / E_\gamma$ , must also have a narrow distribution, since both  $\epsilon_\gamma$  and  $E_K / E_\gamma$  have a narrow spread.

#### 3.2. Isotropic-Equivalent Energies and Beaming Corrections

While there is a clear correlation between  $E_{\gamma,\text{iso}}$  and  $E_{K,\text{iso}}$ , the overall spread in isotropic-equivalent energies appears to be wider than for the beaming-corrected energies of long GRBs, *unless the outflows of short GRBs with the highest energies are strongly collimated*. To date, only GRB 051221a exhibits evidence for significant beaming, with  $f_b \equiv [1 - \cos(\theta_j)] \approx 7.5 \times 10^{-3}$  (Burrows et al. 2006; Soderberg et al. 2006a), while GRB 050709 appears

to have a wide jet with  $f_b \approx 0.06$  (Fox et al. 2005). In general short GRB jets appear to be wider than those of long GRBs with  $\theta_j \gtrsim 10^\circ$  compared to  $\langle \theta_j \rangle \approx 5^\circ$  (Soderberg et al. 2006a).

In the larger sample presented here two additional bursts with known redshifts (061006 and 061210) exhibit steep decays,  $\alpha_X \sim -2$ , at late time, reminiscent of a post jet break evolution (for which the expected value is  $\alpha_X = -p$  with  $p \gtrsim 2$ ; Sari et al. 1999). In addition to GRB 051221a these bursts also have the largest secure values of  $E_{\gamma,\text{iso}}$ . In the case of GRB 061006 there is a clear break at  $t \approx 1 \times 10^5$  s, while for GRB 061210 the light curve is already in the rapid decay phase at the time of the first observations,  $t \approx 2 \times 10^5$  s. Using the conversion from jet break time to opening angle (Sari et al. 1999):

$$\theta_j = 0.21 t_{j,1}^{3/8} (1+z)^{-3/8} E_{\gamma,\text{iso},51}^{-1/8} (\epsilon_\gamma/0.8)^{1/8} n_0^{1/8}, \quad (3)$$

I find  $\theta_j \approx 0.12$  for GRB 061006 and  $\theta_j \lesssim 0.16$  for GRB 061210 ( $n_0 = 10^{-2} \text{ cm}^{-3}$ ). Thus, in the context of jet breaks, the beaming-corrected energies are  $E_\gamma \approx 5 \times 10^{48}$  erg and  $\lesssim 6 \times 10^{48}$  erg, respectively. These values are within a factor of few of  $E_\gamma \approx (1 - 2) \times 10^{49}$  erg inferred for GRB 051221a (Soderberg et al. 2006a), and  $E_\gamma \approx 2.1 \times 10^{48}$  inferred for GRB 050709 (Fox et al. 2005).

If the observed steep decays are indeed due to jets, this leads to a similar inverse correlation between  $E_{\text{iso}}$  and beaming correction that is observed in long GRBs (Frail et al. 2001; Berger et al. 2003a; Bloom et al. 2003), and thus to a narrow distribution of  $E_{\text{rel}} \sim 10^{48} - 10^{49}$  erg. Models of the relativistic outflows from black hole accretion systems relevant for short GRBs indicate  $E_{\text{rel,iso}} \sim 10^{49} - 10^{52}$  with opening angles of  $\sim 5 - 15^\circ$  (Janka et al. 2006), in good agreement with the observations presented here. It is thus possible that short GRBs also exhibit a nearly standard energy release, with an overall scale that is two orders of magnitude smaller than that of long GRBs.

The proposed beaming corrections also impact the nature of the energy extraction mechanism. In particular, extraction via  $\nu\bar{\nu}$  annihilation has been argued to provide at most  $E_{\text{rel}} \sim \text{few} \times 10^{48}$  erg (Rosswog & Ramirez-Ruiz 2002), which may be sufficient if the beaming corrections are valid. On the other hand, if the observed steep decays are not due to jets, then the required large energies likely point to MHD processes such as the Blandford-Znajek mechanism (Blandford & Znajek 1977; Rosswog et al. 2003; Soderberg et al. 2006a), which can produce luminosities in excess of  $10^{52} \text{ erg s}^{-2}$ . These possibilities have to be assessed with a larger sample of bursts, with particular attention to multi-wavelength observations that can differentiate between bona-fide jet breaks and other scenarios (e.g., energy or density variations).

#### 3.3. Correlations with Duration and Host Galaxy Type?

I further investigate whether there is a correlation between the energy release and burst duration ( $T_{90}$ ); Figure 1. For the full sample I find a Spearman's rank correlation coefficient,  $\rho = 0.58$ , or a null hypothesis probability of only  $7.5 \times 10^{-3}$ . Removing GRB 050509b, which has

the shortest duration and smallest fluence, I find  $\rho = 0.49$ , or a probability of  $2.6 \times 10^{-2}$  that the null hypothesis is satisfied. Thus, I conclude that with the present sample the correlation between fluence and duration is suggestive, though not conclusive.

A more physically meaningful test is whether  $E_{\gamma, \text{iso}}$  is correlated with the rest-frame duration,  $T_{90}/(1+z)$ . In this case, I find that for the eight bursts with secure redshifts the null hypothesis has a probability of  $4.5 \times 10^{-2}$ , so an overall correlation does not appear to be statistically meaningful. Curiously, the sample of six bursts with secure redshifts and late-type host galaxies has  $\rho = 0.94$  (null hypothesis probability of only  $6 \times 10^{-4}$ ), but this result is likely due to small number statistics. To conclude, there is some tentative evidence for a positive correlation between energy release and duration, but this will have to be re-assessed with a larger sample.

Similarly, I check for a relation between energy release and host galaxy type. This is the case for Type Ia supernovae (SNe Ia), which tend to have lower luminosities in early-type galaxies than in late-type galaxies, likely due to a dependence on the progenitor ages (Gallagher et al. 2005; Sullivan et al. 2006). Short GRBs also occur in both types of galaxies so a similar trend may shed light on the progenitor properties. In the sample presented here, three bursts have been localized to early-type galaxies (050509b, 050724, and 050813), while six have secure late-type hosts (050709, 051221a, 060801, 061006, 061210, and 061217). As can be seen from Table 1 both groups appear to span the full range of  $\gamma$ -ray energies and X-ray luminosities, suggesting no clear correlation with host galaxy type for the present sample.

#### 3.4. The Nature of the Outliers

Finally, I return to the 20% of outliers with an unusually low ratio of  $F_{X,1}/F_{\gamma}$ . The possible nature of these objects has been discussed by Nakar (2007) who noted several possibilities for the suppressed X-ray flux, in particular a low circumburst density ( $n \lesssim 10^{-5} \text{ cm}^{-3}$ ), typical of the intergalactic medium. This is argued to support the idea of large kick velocities,  $\gtrsim 10^2 \text{ km s}^{-1}$ , for some progenitors. Here I simply note that while all four outliers have fluences at the low end of the distribution,  $F_{\gamma} \lesssim 10^{-7} \text{ erg cm}^{-2}$  (Figure 1), the two with redshift constraints (050813 and 051210) appear to reside at  $z \gtrsim 1.5$  and therefore their energies are at the high end of the overall distribution. No other parameters clearly distinguish the outliers from the bulk of the population that has a narrow distribution of  $\epsilon_{\gamma}$ , suggesting that indeed an extrinsic parameter such as the density is responsible for their low X-ray fluxes.

In the context of a low density interpretation for these outliers, and given the clearly bimodal distribution, I propose the following intriguing possibility. The observed fraction of 20% is similar to predictions of the fraction of short GRBs that may arise from dynamically-formed neutron star binaries in globular clusters (10–30%; Grindlay

et al. 2006). This scenario may naturally explain the required low densities since the intra-cluster medium of globular clusters has a typical limit<sup>6</sup> of  $n \lesssim \text{few} \times 10^{-5} \text{ cm}^{-3}$  (van Loon et al. 2006), likely as a result of gas stripping from frequent passages through the Galactic disk. This scenario also naturally explains the bimodal distribution of  $F_X/F_{\gamma}$  (and hence densities), since the bursts occur in either interstellar environments with  $n \gtrsim 0.05 \text{ cm}^{-3}$  (§3.1) or in the low density environments of globular clusters. Thus, this alternative scenario removes the need for speculative large kick velocities for the outliers (Nakar 2007), which in any case predicts a more uniform distribution of densities and hence  $F_X/F_{\gamma}$  than currently observed.

#### 4. SUMMARY

Based on prompt  $\gamma$ -ray and X-ray afterglow observations of a sample of 17 short GRBs with rapid XRT follow-up I find the following results:

1. The majority of short GRBs (80%) follow a linear correlation between  $E_{\gamma}$  and  $E_K$ , which indicates a narrow distribution of  $\gamma$ -ray efficiency (0.14 dex) with  $\langle \epsilon_{\gamma} \rangle \sim 0.85$ , similar to the values for long GRBs.
2. The observed correlation also indicates that generally  $\nu_X > \nu_c$  and therefore  $n \gtrsim 0.05 \text{ cm}^{-3}$ , indicative of interstellar explosion sites.
3. The isotropic-equivalent energies span  $10^{48} - 10^{52} \text{ erg}$ , similar in width to the distribution for long GRBs, but lower by about two orders of magnitude.
4. Possible beaming corrections for several short bursts with the highest values of  $E_{\gamma, \text{iso}}$  may point to an inverse correlation such that the true energy release has a narrow distribution of  $\sim 10^{48} - 10^{49}$ .
5. There is a possible weak correlation between  $E_{\gamma, \text{iso}}$  and burst duration; no clear relation is evident between energy release and host galaxy type.
6. A small fraction of outliers (20%), with  $F_X/F_{\gamma}$  lower by three orders of magnitude, are likely the result of low circumburst densities. The low density, strong bimodality of the distribution, and similarity in predicted rates, are naturally explained in a globular cluster origin as opposed to large kick velocities.

With these properties of short GRBs well-established the next step is to assess the importance of beaming corrections, and hence the true distribution of total relativistic output from the central engine.

I thank Alicia Soderberg and Ehud Nakar for valuable discussion, and acknowledge support by NASA through Hubble Fellowship grant HST-01171.01 awarded by the Space Telescope Science Institute, which is operated by AURA, Inc. for NASA under contract NAS 5-26555.

#### References

<sup>6</sup>However, a substantial intra-cluster density,  $n \sim 0.07 \text{ cm}^{-3}$ , has been claimed for the globular cluster 47 Tucanae (Freire et al. 2001).

- Barthelmy, S., Gehrels, N., Norris, J., & Sakamoto, T. 2005a, GRB Coordinates Network, 4401
- Barthelmy, S. D., et al. 2005b, *Nature*, 438, 994
- Beardmore, A., Burrows, D. N., Page, K., Marshall, F., & Chester, M. 2005a, GRB Coordinates Network, 4402
- Beardmore, A. P., Page, K. L., Gehrels, N., Greiner, J., Kennea, J., Nousek, J., Osborne, J. P., & Tagliaferri, G. 2005b, GRB Coordinates Network, 4043
- Berger, E. 2006a, GRB Coordinates Network, 5952
- Berger, E. 2006b, in AIP Conf. Proc. 836: Gamma-Ray Bursts in the Swift Era, ed. S. S. Holt, N. Gehrels, & J. A. Nousek, 33
- Berger, E. 2007, GRB Coordinates Network, 5995
- Berger, E., et al. 2006, ArXiv Astrophysics e-prints; astro-ph/0611128
- Berger, E., et al. 2005a, *ApJ*, 634, 501
- Berger, E., Kulkarni, S. R., & Frail, D. A. 2003a, *ApJ*, 590, 379
- Berger, E., et al. 2003b, *Nature*, 426, 154
- Berger, E., et al. 2005b, *Nature*, 438, 988
- Berger, E., et al. 2000, *ApJ*, 545, 56
- Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433
- Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, *ApJ*, 594, 674
- Bloom, J. S., et al. 2007, *ApJ*, 654, 878
- Bloom, J. S., et al. 2006, *ApJ*, 638, 354
- Bloom, J. S. 2006, GRB Coordinates Network, 5880
- Burrows, D. N., et al. 2006, *ApJ*, 653, 468
- Cannizzo, J. K., et al. 2006, *GCNR*, 20
- de Ugarte Postigo, A., et al. 2006, *ApJ*, 648, L83
- Donaghy, T. Q., et al. 2006, ArXiv Astrophysics e-prints; astro-ph/0605570
- Fox, D. B., et al. 2005, *Nature*, 437, 845
- Frail, D. A., et al. 2001, *ApJ*, 562, L55
- Freedman, D. L., & Waxman, E. 2001, *ApJ*, 547, 922
- Freire, P. C., Kramer, M., Lyne, A. G., Camilo, F., Manchester, R. N., & D'Amico, N. 2001, *ApJ*, 557, L105
- Gallagher, J. S., Garnavich, P. M., Berlind, P., Challis, P., Jha, S., & Kirshner, R. P. 2005, *ApJ*, 634, 210
- Gehrels, N., et al. 2005, *Nature*, 437, 851
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'Shin, V., Frederiks, D., & Cline, T. 2005, GRB Coordinates Network, 4394
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'Shin, V., Frederiks, D., & Cline, T. 2006a, GRB Coordinates Network, 4881
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'Shin, V., Frederiks, D., & Cline, T. 2006b, GRB Coordinates Network, 5890
- Golenetskii, S., Aptekar, R., Mazets, E., Pal'shin, V., Frederiks, D., & Cline, T. 2006, GRB Coordinates Network, 5710
- Granot, J., & Sari, R. 2002, *ApJ*, 568, 820
- Grindlay, J., Portegies Zwart, S., & McMillan, S. 2006, *Nature Physics*, 2, 116
- Grupe, D., et al. 2006, *ApJ*, 653, 462

- Hjorth, J., et al. 2005, *Nature*, 437, 859
- Hullinger, D., et al. 2005, GRB Coordinates Network, 4400
- Janka, H.-T., Aloy, M.-A., Mazzali, P. A., & Pian, E. 2006, *ApJ*, 645, 1305
- Kumar, P. 2000, *ApJ*, 538, L125
- La Parola, V., et al. 2006, *A&A*, 454, 753
- Levan, A. J., et al. 2006, *ApJ*, 648, L9
- Markwardt, C., et al. 2005, GRB Coordinates Network, 4037
- Marshall, F., Perri, M., Stratta, G., Barthelmy, S. D., Krimm, H., Burrows, D. N., Roming, P., & Gehrels, N. 2006, GCNR, 18
- Nakar, E. 2007, ArXiv Astrophysics e-prints; astro-ph/0701748
- Pagani, C., La Parola, V., & Burrows, D. N. 2005, GRB Coordinates Network, 3934
- Panaitescu, A., & Kumar, P. 2002, *ApJ*, 571, 779
- Parsons, A., et al. 2005, GRB Coordinates Network, 3935
- Racusin, J. L., Grupe, D., Morris, D., & Strohm, M. 2006, GRB Coordinates Network, 5382
- Roming, P. W. A., et al. 2006, *ApJ*, 651, 985
- Rosswog, S., & Ramirez-Ruiz, E. 2002, *MNRAS*, 336, L7
- Rosswog, S., Ramirez-Ruiz, E., & Davies, M. B. 2003, *MNRAS*, 345, 1077
- Sari, R., Piran, T., & Halpern, J. P. 1999, *ApJ*, 519, L17
- Sato, G., et al. 2005, GRB Coordinates Network, 3793
- Sato, G., et al. 2006a, GRB Coordinates Network, 5064
- Sato, G., et al. 2006b, GRB Coordinates Network, 5381
- Schady, P., et al. 2006, GCNR, 6
- Soderberg, A. M., et al. 2006a, *ApJ*, 650, 261
- Soderberg, A. M., et al. 2004, *Nature*, 430, 648
- Soderberg, A. M., et al. 2006b, *Nature*, 442, 1014
- Sullivan, M., et al. 2006, *ApJ*, 648, 868
- Troja, E., Burrows, D. N., & Gehrels, N. 2006, GRB Coordinates Network, 5093
- Urata, Y., et al. 2006, GRB Coordinates Network, 5917
- van Loon, J. T., Stanimirović, S., Evans, A., & Muller, E. 2006, *MNRAS*, 365, 1277
- Villasenor, J. S., et al. 2005, *Nature*, 437, 855
- Yost, S. A., Harrison, F. A., Sari, R., & Frail, D. A. 2003, *ApJ*, 597, 459
- Ziaeeepour, H., et al. 2006, GCNR, 21

TABLE 1  
 $\gamma$ -RAY AND X-RAY PROPERTIES OF SHORT GRBs

| GRB     | $z$                     | $T_{90}$<br>(s)   | $F_{\gamma}^a$<br>(erg cm $^{-2}$ ) | $\Delta t$<br>(s) | $F_X$<br>(erg cm $^{-2}$ s $^{-1}$ ) | $\alpha_X$ | Refs.       |
|---------|-------------------------|-------------------|-------------------------------------|-------------------|--------------------------------------|------------|-------------|
| 050509b | 0.226                   | $0.040 \pm 0.004$ | $(9.5 \pm 2.5) \times 10^{-9}$      | 200               | $(3.6 \pm 1.1) \times 10^{-13}$      | -1.1       | 1–2         |
| 050709  | 0.1606                  | $0.070 \pm 0.010$ | $(3.0 \pm 0.4) \times 10^{-7}{}^b$  | $2.2 \times 10^5$ | $(3.5 \pm 0.5) \times 10^{-15}$      | -1.0       | 3–5         |
| 050724  | 0.257                   | $3.000 \pm 1.000$ | $(3.9 \pm 1.0) \times 10^{-7}$      | 6625              | $(8.8 \pm 2.1) \times 10^{-13}$      | -1.0       | 6–8         |
| 050813  | $\sim 1.8$              | $0.600 \pm 0.100$ | $(1.2 \pm 0.5) \times 10^{-7}$      | 2100              | $(3.0 \pm 1.2) \times 10^{-14}$      | -2.0       | 9–11        |
| 050906  | ...                     | $0.128 \pm 0.016$ | $(5.9 \pm 3.2) \times 10^{-8}$      | 79                | $< 8.0 \times 10^{-14}$              | ...        | 12–13       |
| 050925  | ...                     | $0.068 \pm 0.027$ | $(7.5 \pm 0.9) \times 10^{-8}$      | 100               | $< 3.0 \times 10^{-14}$              | ...        | 14–15       |
| 051210  | $> 1.55$                | $1.200 \pm 0.200$ | $(8.1 \pm 1.4) \times 10^{-8}$      | 840               | $(2.4 \pm 0.6) \times 10^{-12}$      | -2.6       | 16–17       |
| 051221a | 0.5465                  | $1.400 \pm 0.200$ | $(1.2 \pm 0.1) \times 10^{-6}{}^c$  | $3.9 \times 10^4$ | $(1.2 \pm 0.1) \times 10^{-12}$      | -1.2       | 18–19       |
| 051227  | ...                     | $8.000 \pm 0.200$ | $(2.3 \pm 0.3) \times 10^{-7}$      | $3.5 \times 10^4$ | $(1.8 \pm 0.6) \times 10^{-13}$      | -1.1       | 20–22       |
| 060121  | ...                     | $1.970 \pm 0.060$ | $(4.8 \pm 0.3) \times 10^{-6}{}^d$  | $2.2 \times 10^4$ | $(2.1 \pm 0.5) \times 10^{-12}$      | -1.2       | 23–25       |
| 060313  | $< 1.7$                 | $0.700 \pm 0.100$ | $(1.1 \pm 0.1) \times 10^{-6}{}^e$  | $1.0 \times 10^5$ | $(1.5 \pm 0.3) \times 10^{-13}$      | -1.5       | 26          |
| 060502b | ... <sup>f</sup>        | $0.090 \pm 0.020$ | $(4.0 \pm 0.5) \times 10^{-8}$      | $2.8 \times 10^4$ | $(2.4 \pm 1.2) \times 10^{-14}$      | -1.1       | 10,17,27–29 |
| 060801  | 1.1304                  | $0.500 \pm 0.100$ | $(8.1 \pm 1.0) \times 10^{-8}$      | $7.0 \times 10^4$ | $< 3.6 \times 10^{-14}$              | ...        | 10,17,30–31 |
| 061006  | 0.4377                  | 0.420             | $(1.4 \pm 0.1) \times 10^{-6}{}^g$  | $1.0 \times 10^5$ | $(1.0 \pm 0.4) \times 10^{-13}$      | -1.9       | 17,32       |
| 061201  | $\sim 0.1$ <sup>h</sup> | $0.800 \pm 0.100$ | $(3.3 \pm 0.3) \times 10^{-7}{}^i$  | $3.4 \times 10^4$ | $(2.0 \pm 0.5) \times 10^{-13}$      | -1.9       | 33–35       |
| 061210  | 0.4095                  | 0.190             | $(1.1 \pm 0.2) \times 10^{-6}{}^j$  | $2.3 \times 10^5$ | $(1.4 \pm 0.5) \times 10^{-13}$      | -2.0       | 17,36       |
| 061217  | 0.8270                  | $0.212 \pm 0.041$ | $(4.6 \pm 0.8) \times 10^{-8}$      | 1340              | $(4.7 \pm 1.0) \times 10^{-13}$      | -0.6       | 17,37       |

NOTE.—Prompt emission and X-ray Properties of the short GRB discussed in this paper, including (i) GRB name, (ii) redshift, (iii) duration, (iv)  $\gamma$ -ray fluence, (v) time of X-ray observations, (vi) X-ray flux, (vii) X-ray temporal decay rate, and (viii) references.

<sup>a</sup> Fluence is in the 15 – 150 keV energy band unless otherwise noted.

<sup>b</sup> The fluence is in the 30 – 400 keV band and for the initial short pulse only; the total fluence including the  $\sim 130$  s duration soft tail is  $F_{\gamma} \approx 1.5 \times 10^{-6}$  erg cm $^{-2}$  in the 2 – 400 keV band (Villasenor et al. 2005).

<sup>c</sup> The fluence in the 20 – 2000 keV band is  $3.2 \times 10^{-6}$  erg cm $^{-2}$  (Golenetskii et al. 2005).

<sup>d</sup> The fluence is in the 2 – 400 keV band.

<sup>e</sup> The fluence in the 20 – 2000 keV band is  $1.4 \times 10^{-5}$  erg cm $^{-2}$  (Golenetskii et al. 2006a).

<sup>f</sup> A redshift of  $z = 0.257$  has been proposed by Bloom et al. (2007) based on a galaxy located 17.5'' away from the XRT position, but a fainter galaxy of unknown redshift has been identified within the error circle (Berger et al. 2006).

<sup>g</sup> The fluence in the 20 – 2000 keV band is  $3.6 \times 10^{-6}$  erg cm $^{-2}$  (Golenetskii et al. 2006).

<sup>h</sup> The nature of the host is not clear, given the location of an Abell cluster at  $z = 0.0865$  about 8' away (Bloom 2006; Berger 2007), as well as a galaxy at  $z = 0.111$  about 17'' away (Berger 2006a).

<sup>i</sup> The fluence in the 20 – 3000 keV band is  $5.3 \times 10^{-6}$  erg cm $^{-2}$  (Golenetskii et al. 2006b).

<sup>j</sup> The fluence in the 100 – 1000 keV band is  $2.0 \times 10^{-6}$  erg cm $^{-2}$  (Urata et al. 2006).

References: [1] Gehrels et al. (2005); [2] Bloom et al. (2007); [3] Villasenor et al. (2005); [4] Fox et al. (2005); [5] Hjorth et al. (2005); [6] Barthelmy et al. (2005b); [7] Berger et al. (2005b); [8] Grupe et al. (2006); [9] Sato et al. (2005); [10] Nakar (2007); [11] Berger (2006b); [12] Pagani et al. (2005); [13] Parsons et al. (2005); [14] Markwardt et al. (2005); [15] Beardmore et al. (2005b); [16] La Parola et al. (2006); [17] Berger et al. (2006); [18] Burrows et al. (2006); [19] Soderberg et al. (2006a); [20] Hullinger et al. (2005); [21] Barthelmy et al. (2005a); [22] Beardmore et al. (2005a); [23] Donaghy et al. (2006); [24] Levan et al. (2006); [25] de Ugarte Postigo et al. (2006); [26] Roming et al. (2006); [27] Sato et al. (2006a); [28] Troja et al. (2006); [29] Bloom et al. (2007); [30] Sato et al. (2006b); [31] Racusin et al. (2006); [32] Schady et al. (2006); [33] Marshall et al. (2006); [34] Berger (2006a); [35] Berger (2007); [36] Cannizzo et al. (2006); [37] Ziaeeepour et al. (2006).




FIG. 1.— X-ray flux normalized to  $t = 1$  d and burst duration ( $T_{90}$ ) plotted against the  $\gamma$ -ray fluence for the complete sample of 17 short GRBs with prompt XRT follow-up. The majority of the bursts (80%) follow a clear linear relation between  $F_{X,1}$  and  $F_\gamma$  suggesting a relatively standard fraction of energy emitted in  $\gamma$ -rays. At the same time, a subset of the short bursts have X-ray fluxes that are suppressed by about three orders of magnitude. The projected histograms indicate the full range of values for each observed quantity (arrows designate upper limits on the X-ray flux). There is also a possible correlation between  $T_{90}$  and  $F_\gamma$  (with a large dispersion) – the null hypothesis of no correlation has a probability of only 0.75%.






FIG. 2.— X-ray luminosity normalized to  $t = 1$  d (a proxy for  $\epsilon_e E_{K,iso}$ ) plotted against  $E_{\gamma,iso}$  for the short bursts with a known redshift (solid black circles), redshift constraints (open black circles) and without any redshift information (gray symbols connected by dotted lines). For the latter I plot the inferred values at  $z = 0.1$  and  $z = 1$ , corresponding roughly to the lowest and highest redshifts securely measured to date. The isotropic-equivalent relativistic energies are as least as high as  $10^{51}$  erg, and may approach  $\text{few} \times 10^{52}$  erg for some short bursts (see also Berger et al. 2006).